

Table 9. Estimated numbers of persons living with HIV/AIDS at the end of 2004, by race/ethnicity, sex, and transmission category—35 areas with confidential name-based HIV infection reporting

Transmission category	White, not Hispanic		Black, not Hispanic		Hispanic		Asian/Pacific Islander		American Indian/Alaska Native		Total ^a	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	101,119	76	63,112	46	31,412	54	1,415	66	879	61	199,085	60
Injection drug use	12,750	10	33,237	24	14,990	26	267	12	218	15	61,799	19
Male-to-male sexual contact and injection drug use	10,667	8	9,030	7	3,237	6	74	3	191	13	23,337	7
Heterosexual contact	6,260	5	29,458	22	8,109	14	334	16	146	10	44,655	13
Other ^b	1,580	1	1,617	1	411	1	50	2	18	1	3,702	1
Subtotal	132,376	100	136,454	100	58,159	100	2,141	100	1,452	100	332,578	100
Female adult or adolescent												
Injection drug use	8,104	34	19,554	25	5,521	30	96	16	168	32	33,621	27
Heterosexual contact	15,256	64	57,991	73	12,636	68	470	79	347	66	87,262	71
Other ^b	481	2	1,626	2	357	2	30	5	10	2	2,523	2
Subtotal	23,841	100	79,170	100	18,513	100	596	100	525	100	123,405	100
Child (<13 yrs at diagnosis)												
Perinatal	793	83	4,013	91	1,228	90	23	85	18	95	6,100	90
Other ^c	161	17	389	9	138	10	4	15	1	5	704	10
Subtotal	953	100	4,402	100	1,366	100	27	100	19	100	6,804	100
Total^d	157,172		220,028		78,039		2,765		1,996		462,792	

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

Data include persons with a diagnosis of HIV infection. This includes persons with a diagnosis of HIV infection only, a diagnosis of HIV infection and a later AIDS diagnosis, and concurrent diagnoses of HIV infection and AIDS.

Since 2000, the following 35 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, Guam and the U.S. Virgin Islands. Since July 1997, Florida has had confidential name-based HIV infection reporting only for new diagnoses.

^a Includes 2,793 persons of unknown race or multiple races.

^b Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

^c Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^d Includes 5 persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.